









## Crop Weather Calendar of Rabi Chili: Rajshahi Region (Districts:Rajshahi, Chapainawabgonj, Naoga, Natore) Bangladesh

Rajshahi Region	Rabi Chili										Life span: 170-180 days											
Months	November					December					January				February				March			
Std.Week/Normal	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8	9	10	11	12	13
Rainfall (mm)	7.0	5.0	1.0	0.0	2.0	1.0	3.0	0.0	4.0	1.0	1.0	2.0	1.0	4.0	3.0	2.0	3.0	5.0	4.0	3.0	4.0	14.0
Max. Temp. (°C)	31.7	31.0	30.4	30.4	29.6	27.0	26.0	25.4	24.4	24.0	23.9	24.0	24.9	25.6	26.7	28.1	29.3	30.4	32.1	33.1	34.6	35.0
Min. Temp. (°C)	23.1	21.4	20.7	19.3	17.6	14.0	13.1	12.4	11.6	11.0	10.9	10.9	10.9	12.0	12.3	13.6	14.1	15.1	16.4	17.6	19.4	20.6
Mean Temp. (°C)	27.4	26.2	25.6	24.9	23.6	20.5	19.6	18.9	18.0	17.5	17.4	17.5	17.9	18.8	19.5	20.9	21.7	22.8	24.3	25.4	27.0	27.8
RHmax (%)	96.1	96.0	95.9	96.1	95.7	94.7	95.9	95.3	95.7	95.6	95.1	95.3	95.3	95.3	94.0	93.9	92.0	91.6	90.3	88.7	88.3	89.1
RHMin (%)	60.9	57.4	55.9	51.9	49.4	46.9	50.1	50.0	50.0	50.1	48.4	47.9	45.7	44.7	42.0	40.3	36.6	34.6	32.7	30.4	30.4	35.9
RHmean (%)	78.5	76.7	75.9	74.0	72.6	70.8	73.0	72.7	72.9	72.9	71.8	71.6	70.5	70.0	68.0	67.1	64.3	63.1	61.5	59.4	59.4	62.5
SShr (hrs)	53.0	56.0	55.0	57.0	54.0	51.0	47.0	46.0	44.0	42.0	42.0	46.0	52.0	51.0	55.0	56.0	58.0	59.0	59.0	59.0	59.0	52.0
WD (deg)	146	146	122	137	151	143	146	144	167	146	161	174	196	198	195	197	215	209	220	214	208	186
WS(Km/hr)	3.9	4.3	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.9	3.7	3.7	4.4	3.9	5.0	3.9	5.0	5.0	5.4	5.4	5.6	5.6
																						
Stage	Spouting & Seedling					vegetative growth					Flowering & fruiting					Harvesting						
<b>Favorable Environment</b>																						
Temperature	The ideal average monthly temperatures range between 21°C and 30 °C, with an average of 18 °C; Germination and growth decreases at lower temperature (10°C); Fruit set is adversely affected at temperatures above 35 °C as well as wilting of young parts, and slow growth occurred, especially with low air humidity or dry winds.																					
Relative Humidity	Sensitive to excess moisture and the high relative humidity; The optimum relative humidity varies from 50% to 70%.																					
Soil Temperature	Chili seeds can germinate and grow normally at soil temperatures of 15°C to 25°C, with the optimum around 20°C. Chili can tolerate moderately cold soil temperatures of 13°C to 15°C.																					
Sunshine Hour	At least 4 to 6 hours direct sunlight per day																					
<b>Favorable environmental conditions for insects and diseases</b>																						
Mites	Maximum temp.: 33 to 35°C; Raise a generation in 1 week under optimal conditions (25°C and high relative humidity); Minimum Temp.: 12 to 14°C;The prevailing high temperature and low relative humidity in absence of rainfall favored the population development of mites to reach the peak abundance.																					
Aphids	Moderate to high temperature and relative humidity favors population buildup of aphids																					
Thrips	Grow minimum temperature 9.7°C and maximum temperatures, 33.0°C; Population remains more abundant during prolonged dry conditions than in moist rainy periods; Heavy rainfall reduces population																					
White fly	Population increases at high temperatures; white fly population is low at humidity ≤45% and ≥85% and abundant at 70–85% RH																					
Fruit Borer	The upper development threshold temperature for all stages is 37 °C, and 40 °C is lethal; The optimum temperatures for the development of eggs, larvae, and pupae are around 24–27 °C; RH: 85% to ≤35%.																					
Leaf curl	Vector (White fly) population and virus transmission increased at 35 °C (max. temp.); Disease development correlated with relative humidity as the humidity decreased the disease incident decreased; Disease development favored by certain rainfall																					
Die back/Anthracnose	Max. Temp.: 28 °C to 32°C for disease development; Mini. Temp.: 22°C to 25°C for disease development; Initial infection required 95% or above relative humidity for 24 hrs; Frequent rainy and warm weather alone with prolonged leaf wetness favored disease development.																					
<b>Weather Warning</b>																						
Rainfall	Early rainfall (10 to 15 mm per day)																					
Cloudy weather	Cloudy weather																					
Drought	Short to moderate drought (7 to 10 days)																					
Duration of staying wet	2 to 3 days																					

Hailstorm			Hailstorms with early rainfall
Windspeed			≤20 Km/hr
Temperature	Germination and growth decreases at lower temperature (10°C), Fruit set is adversely affected at temperatures above 35 °C		

### Crop Weather Calendar of Kharif Chili: Rajshahi Region (Districts:Rajshahi, Chapainawabgonj, Naoga, Natore) Bangladesh

Rajshahi Region	Kharif Chili													Life span: 170-180 days								
months	April				may					June				July				August				
Months	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35
Std.Week/Normal	13.0	10.0	19.0	17.0	24.0	29.0	33.0	36.0	34.0	48.0	51.0	78.0	72.0	66.0	71.0	75.0	72.0	52.0	57.0	64.0	53.0	57.0
Rainfall (mm)	36.1	36.7	36.4	35.9	35.4	35.3	35.1	35.0	35.3	35.0	33.9	33.1	33.0	32.7	32.9	32.3	32.6	32.9	32.9	32.7	33.0	33.0
Max. Temp. (°C)	22.1	23.0	23.4	23.9	23.7	24.3	24.6	25.0	25.4	25.9	26.0	26.0	26.0	26.0	26.0	26.0	26.0	26.0	26.1	26.0	26.1	26.0
Min. Temp. (°C)	29.1	29.9	29.9	29.9	29.6	29.8	29.9	30.0	30.4	30.5	30.0	29.6	29.5	29.4	29.5	29.2	29.3	29.5	29.5	29.4	29.6	29.5
Mean Temp. (°C)	88.9	89.0	90.1	90.7	92.1	92.7	93.3	93.1	93.7	94.6	95.4	96.1	96.6	96.6	96.9	97.0	96.9	96.7	96.6	96.7	96.1	96.4
RHmax (%)	34.3	34.6	41.4	45.1	50.0	51.9	55.6	58.3	58.9	62.6	67.9	71.1	73.4	73.7	74.3	75.3	73.9	72.6	72.6	74.1	72.0	70.9
RHMin (%)	61.6	61.8	65.8	67.9	71.1	72.3	74.5	75.7	76.3	78.6	81.7	83.6	85.0	85.2	85.6	86.2	85.4	84.7	84.6	85.4	84.1	83.7
RHmean (%)	57.0	57.0	57.0	56.0	52.0	53.0	54.0	48.0	51.0	48.0	35.0	31.0	31.0	30.0	30.0	30.0	34.0	34.0	36.0	33.0	34.0	40.0
SShr (hrs)	185	185	267	147	136	136	135	126	133	139	129	127	132	141	134	138	140	131	131	141	143	133
WD (deg)	6.1	6.1	8.2	7.4	7.4	7.2	7.2	7.2	7.2	6.7	6.7	6.8	6.8	5.7	6.1	6.3	6.7	6.3	5.6	5.7	5.6	5.6
																						
Stage	Spouting & Seedling				vegetative growth					Flowering & fruiting				Harvesting								
<b>Favorable Environment</b>																						
Temperature	The ideal average monthly temperatures range between 21°C and 30 °C, with an average of 18 °C; Germination and growth decreases at lower temperature (10°C); Fruit set is adversely affected at temperatures above 35 °C as well as wilting of young parts, and slow growth occurred, especially with low air humidity or dry winds.																					
Relative Humidity	Sensitive to excess moisture and the high relative humidity; The optimum relative humidity varies from 50% to 70%.																					
Soil Temperature	Chili seeds can germinate and grow normally at soil temperatures of 15°C to 25°C, with the optimum around 20°C. Chili can tolerate moderately cold soil temperatures of 13°C to 15°C.																					
Sunshine Hour	At least 4 to 6 hours direct sunlight per day																					
<b>Favorable environmental conditions for insects and diseases</b>																						
Mites	Maximum temp.: 33 to 35°C; Raise a generation in 1 week under optimal conditions (25°C and high relative humidity); Minimum Temp.: 12 to 14°C;The prevailing high temperature and low relative humidity in absence of rainfall favored the population development of mites to reach the peak abundance.																					
Aphids	Moderate to high temperature and relative humidity favors population buildup of aphids																					
Thrips	Grow minimum temperature 9.7°C and maximum temperatures, 33.0°C; Population remains more abundant during prolonged dry conditions than in moist rainy periods; Heavy rainfall reduces population																					
White fly	Population increases at high temperatures; white fly population is low at humidity ≤45% and ≥85% and abundant at 70–85% RH																					
Fruit Borer	The upper development threshold temperature for all stages is 37 °C, and 40 °C is lethal; The optimum temperatures for the development of eggs, larvae, and pupae are around 24–27 °C; RH: 85% to ≤35%.																					
Leaf curl	Vector (White fly) population and virus transmission increased at 35 °C (max. temp.); Disease development correlated with relative humidity as the humidity decreased the disease incident decreased; Disease development favored by certain rainfall																					
Die back/Anthracnose	Max. Temp.: 28 °C to 32°C for disease development; Mini. Temp.: 22°C to 25°C for disease development; Initial infection required 95% or above relative humidity for 24 hrs;																					

		Frequent rainy and warm weather alone with prolonged leaf wetness favored disease development.
<b>Weather Warning</b>		
Rainfall		Early rainfall (10 to 15 mm per day)
Cloudy weather		Cloudy weather
Drought		Short to moderate drought (7 to 10 days)
Duration of staying wet		2 to 3 days
Hailstorm		Hailstorms with early rainfall
Windspeed		≤20 Km/hr
Temperature		Germination and growth decreases at lower temperature (10°C), Fruit set is adversely affected at temperatures above 35 °C