

























### Crop insect weather calendar of the Rabi season Chili (Fruit borer): Cumilla Region (Districts: Cumilla, Chandpur, B.Barria) Bangladesh

Cumilla Region		<i>Rabi season Chili: Fruit borer insect</i>												Life span: 120-130 days													
Weather warning	Months	October				November				December				January			February			March							
	Max. Temp. (°C)	The upper development threshold temperature for all stages is 37 °C, and 40 °C is lethal; The optimum temperatures for the development of eggs, larvae, and pupae are around 24–27 °C																									
	Min. Temp. (°C)	Threshold for lower temperatures is around 10–12 °C																									
	RH Max (%)	Above 85%, Fecundity is adversely affected by low humidity along with high temperature																									
	RH Min (%)	<35%																									
	Rainfall (mm)	<i>Negative correlation with rainfall</i>																									
Weekly normal weather	Std. Week/Normal	40	41	42	43	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8	9	10	11	12	13
	Rainfall (mm)	51.0	46.5	30.5	16.0	11.5	18.0	8.5	4.0	2.5	0.0	4.5	1.0	1.5	0.0	0.0	0.0	1.5	3.0	3.0	6.0	7.5	4.5	7.0	9.0	19.0	27.0
	Max. Temp. (°C)	32.3	31.9	31.8	31.2	31.0	30.5	29.8	29.0	28.2	27.7	26.5	25.8	25.4	24.7	24.5	25.0	25.4	16.2	27.3	28.1	29.1	29.8	30.9	32.0	32.2	32.5
	Min. Temp. (°C)	25.4	24.7	24.3	22.8	21.8	21.0	19.3	18.1	16.9	15.8	15.0	14.2	13.4	13.0	12.7	12.7	12.6	14.2	14.5	16.2	16.7	18.1	19.2	19.9	21.5	22.0
	RHmax (%)	95.6	95.7	95.5	95.2	95.3	95.6	95.3	95.4	95.8	95.8	96.2	96.0	96.0	96.0	95.5	95.3	94.4	94.6	94.4	93.9	94.1	93.9	93.9	93.5	93.7	94.0
	RHMin (%)	70.2	68.1	65.8	61.9	60.6	58.2	54.8	52.6	52.2	50.6	51.9	52.1	51.3	51.8	51.2	49.1	46.4	47.7	45.2	46.8	46.1	46.8	47.4	47.7	52.6	55.9
SShr (hrs)	39.0	43.0	44.5	50.5	47.0	49.5	50.0	50.5	48.5	47.5	42.5	40.5	40.0	40.0	37.0	43.5	47.0	47.0	50.0	49.5	50.5	52.5	52.5	52.5	51.0	48.5	
Stage																											
		Spouting & Seedling				vegetative growth				Flowering & fruiting						Harvesting											
Insect Life cycle																											
										Eggs (2 to 4 days)		Larva (15 to 18 days)		Pupa (7 to 10 days)		Adults (7 to 10 days)		Eggs (2 to 4 days)		Larva (15 to 18 days)		Pupa (7 to 10 days)		Adults (7 to 10 days)			

### Crop insect weather calendar of the Rabi season Chili: Cumilla Region (Districts: Cumilla, Chandpur, B.Baria) Bangladesh

Cumilla Region		Kharif season Chili: Fruit borer insect															Life span: 170-180 days										
Weather warming	Months	March					April					May					June			July			August				
	Max. Temp. (°C)	The upper development threshold temperature for all stages is 37 °C, and 40 °C is lethal; The optimum temperatures for the development of eggs, larvae, and pupae are around 24–27 °C																									
	Min. Temp. (°C)	Threshold for lower temperatures is around 10–12 °C																									
	RH Max (%)	Above 85%, Fecundity is adversely affected by low humidity along with high temperature																									
	RH Min (%)	<35%																									
	Rainfall (mm)	<i>Negative correlation with rainfall</i>																									
Weekly normal	Std.Week/Normal	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35
	Rainfall (mm)	7.0	9.0	19.0	27.0	25.5	23.0	33.0	42.0	54.5	63.5	66.0	71.0	80.0	74.0	98.5	88.5	97.5	96.0	88.0	107.0	95.0	84.0	66.5	82.5	73.0	66.0
	Max. Temp. (°C)	30.9	32.0	32.2	32.5	32.5	33.0	33.2	33.1	32.9	33.2	33.2	32.9	33.1	32.7	31.9	31.9	31.4	31.2	31.5	31.3	31.5	32.0	32.0	31.9	32.0	32.0
	Min. Temp. (°C)	19.2	19.9	21.5	22.0	23.1	23.5	23.7	23.5	23.7	24.3	24.7	25.2	25.6	25.8	25.7	26.0	26.0	26.0	26.0	25.7	26.0	25.9	26.0	26.0	26.0	25.9
	RHmax (%)	93.9	93.5	93.7	94.0	94.1	93.8	93.9	94.1	94.2	94.0	94.5	94.3	94.6	95.0	95.0	95.2	95.2	95.4	95.0	95.5	95.3	94.9	94.8	94.9	95.3	95.0
	RHMin (%)	47.4	47.7	52.6	55.9	58.7	59.6	60.9	61.7	63.0	63.5	64.8	66.7	69.0	70.9	73.9	75.1	76.8	76.8	75.9	76.6	75.2	74.2	73.9	74.8	74.0	73.4
SShr (hrs)	52.5	52.5	51.0	48.5	51.0	52.0	52.5	51.0	47.5	47.5	46.5	41.0	38.0	36.0	26.5	29.0	26.0	23.5	30.0	26.5	31.0	30.0	31.5	32.0	30.5	34.5	
Stage																											
	Spouting & Seedling					vegetative growth					Flowering & fruiting					Harvesting											
Insect Life cycle																											
	Eggs (2 to 4 days)		Larva (15 to 18 days)					Pupa (7 to 10 days)			Adults (7 to 10 days)			Eggs (2 to 4 days)		Larva (15 to 18 days)			Pupa (7 to 10 days)		Adults (7 to 10 days)						