Crop Disease Weather Calendar of *Rabi* season Pointed gourd (Powdery mildew disease): Bogura region (Districts: Pabna, Sirajganj, Bogura, Joypurhat), Bangladesh

Region: Bogura			Rabi season Pointed Gourd: Powdery mildew disease															Duration: 120-150 days							
ρ0	Months				No	ovember		December				January Feb				Febru	ruary				March				
Weather warning	Max. Temp. (°C)	For infection: 32 °C and for disease development: 27 °C (warm, dry conditions)																							
	Min. Temp. (°C)	For infection: 10 °C and for disease development: 20 °C																							
	RH Max (%)	90% (Two hour or more hours of high humidity or dew on host)																							
	RH Min (%)		50%																						
	Rainfall (mm)				infall also favors the development of the disease																				
al weather	Std. Week/Normal	44	45	46			47	48	49	50	51	52	1	2	3	4	5	6	7	8	9	10	11		13
	Rainfall (mm)	5.0	5.0	0.0			0.0	3.0	0.0	3.0	0.0	4.0	0.0	2.0	2.0	1.0	1.0	1.0	4.0	3.0	5.0	2.0	4.0	3.0	৮.০
	Max. Temp. (°C)	31.0	30.7	30.1			29.7	28.9	27.7	26.3	25.0	25.7	24.0	24.0	24.3	25.4	25.9	26.9	27.7	28.6	29.3	30.9	31.9	32.9	32.9
normal	Min. Temp. (°C)	21.0	19.9	18.3			17.3	15.7	15.0	14.1	12.0	13.4	12.0	11.4	11.4	12.0	12.9	13.6	14.6	15.4	16.4	17.4	18.3	20.0	21.1
Weekly n	RH max (%)	94.3	94.6	94.1			93.5	93.9	93.7	94.6	94.3	94.6	94.1	94.3	94.3	93.9	93.1	92.1	91.4	90.4	88.7	89.1	86.4	86.3	87.7
	RH Min (%)	55.6	52.4	49.9			48.3	47.5	48.1	50.6	50.6	50.0	50.9	49.0	47.7	46.0	14.9	41.4	40.3	39.1	38.0	36.93	37.1	39.1	45.3
>	SS hr (hrs)	52.0	55.0	55.0			56.0	56.0	54.0	48.0	49.0	46.0	44.0	44.0	46.0	50.0	0.0	52.0	53.0	56.0	60.0	59.0	61.0	58.0	56.0
	Growth stage																								
			Spouting & Seedling							vegetative growth Flowering & fr							ing Harvesting								
Disease Life cycle																									
		Spore Spore Germ tube Disease deposition germination penetration development						Disease cycle continued up to fruiting stage																	

Crop Disease Weather Calendar of *Kharif* season Pointed gourd (Powdery mildew disease): Bogura region (Districts: Pabna, Sirajganj, Bogura, Joypurhat), Bangladesh

Region: Bogura			Kharif season Pointed Gourd: Powdery mildew disease														Duration: 100-120 days									
Weather	Months	March							April May								June				July					
	Max. Temp. (°C)	For infection: 32 °C and for disease development: 27 °C (warm, dry conditions)																								
	Min. Temp. (°C)	For infection: 10 °C and for disease development: 20 °C																								
	RH Max (%)	90% (Two hour or more hours of high humidity or dew on host)																								
	RH Min (%)									509	%															
	Rainfall (mm)	Lack of rainfall also favors the development of the disease																								
	Std. Week/Normal	9	10		11		12	13	14	15	16	17	18	19	20	21	22	23	24	25		27	28	29	30	
nal	Rainfall (mm)	5.0	2.0	4.0			3.0	8.0	8.0	12.0	26.0	28.0	37.0		45.0								80.0	91.0		
Weekly normal weather	Max. Temp. (°C)		30.9	31.9			32.9	32.9	33.9	34.3	34.0	33.6	33.1		33.4				_	_			31.9	32.0	32.1	
dy r	Min. Temp. (°C)		17.4		18.3		20.0	21.1	22.3	23.0	23.0	23.1	23.3		24.3								26.1	26.0	26.1	
eekl		88.7			86.4		86.3	87.7	88.0	88.3	90.6		92.4		92.7	_	_	-	-	-	-	_	95.9		95.9	
>	RH Min (%)	_	36.9		37.1		39.1	45.3	44.4	45.9	50.7	54.9	57.3		60.0									74.4	73.3	
	SS hr (hrs)	60.0	59.0		61		58.0	56.0	56.0	56.0	55.0	52.0	48.0	50.0	49.0	46.0	44.0	39.0	33.0	28.0	26.0	28.0	31.0	29.0	28.0	
	Growth stage	Spouting & Seedling																								
									vegetative growth				Flowering & fruiting						Harvesting							
Disease Life cycle																										
		Spore Spore Germ tube Disease deposition germination penetration development							Disease cycle continued up to fruiting stage																	