Crop Insect Weather Calendar of *Rabi* season Pointed gourd (Epilachna beetle insect): Bogura region (Districts: Pabna, Sirajganj, Bogura, Joypurhat)

Region: Bogura			Rabi season Pointed Gourd: Epilachna beetle insect														Duration: 120-150 days							
Weather warning	Months			Novem	ber	December						February				March								
	Max. Temp. (°C)					High temperatu	ıre (35 °C);	25°C temp	erature fa	avored both f	ecundity and foo	d consumpti	ion of the	Epilachn	a beetle as w	vell as larva	l develop	oment						
	Min. Temp. (°C)								Pop	oulation decre	eased with the te	mperature f	all below:	23 ºC										
	RH Max (%)	High relative humidity favored Epilachna beetle (70% or above)																						
	RH Min (%)	30%																						
	Rainfall (mm)	Rainfall is favorable for the activity of Epilachna beetle population																						
er	Std. Week	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8	9	10	11	12	13	
al weather	Rainfall (mm)	5.0	5.0	0.0	0.0	3.0	0.0	3.0	0.0	4.0	0.0	2.0	2.0	1.0	1.0	1.0	4.0	3.0	5.0	2.0	4.0	3.0	৮.০	
	Max. Temp. (°C)	31.0	30.7	30.1	29.7	28.9	27.7	26.3	25.0	25.7	24.0	24.0	24.3	25.4	25.9	26.9	27.7	28.6	29.3	30.9	31.9	32.9	32.9	
norm	Min. Temp. (°C)	21.0	19.9	18.3	17.3	15.7	15.0	14.1	12.0	13.4	12.0	11.4	11.4	12.0	12.9	13.6	14.6	15.4	16.4	17.4	18.3	20.0	21.1	
	RH max (%)	94.3	94.6	94.1	93.5	93.9	93.7	94.6	94.3	94.6	94.1	94.3	94.3	93.9	93.1	92.1	91.4	90.4	88.7	89.1	86.4	86.3	87.7	
Weekly	RH Min (%)	55.6	52.4	49.9	48.3	47.5	48.1	50.6	50.6	50.0	50.9	49.0	47.7	46.0	44.9	41.4	40.3	39.1	38.0	36.93	37.1	39.1	45.3	
>	SS hr (hrs)	52.0	55.0	55.0	56.0	56.0	54.0	48.0	49.0	46.0	44.0	44.0	46.0	50.0	50.0	52.0	53.0	56.0	60.0	59.0	61.0	58.0	56.0	
	Growth stage													海 東 東 東 東 東										
			Spouting	& Seedlin	g	vegetative growth							Harvesting											
	Insect Life cycle									36														
misett zire tytic						Eggs (2 to 5 days)	Grub Pupa Adult (4 to 9 days) (9 to 15 days) (3 to 10 c				Eggs (2 to 5 days)	Grub (4 to 9 days)	(4 to 9 Pupa			Adults Eggs (3 to 10 (2 to 5 days)								

Crop Insect Weather Calendar of *Kharif* season Pointed gourd (Epilachna beetle insect): Bogura region (Districts: Pabna, Sirajganj, Bogura, Joypurhat)

Reg	Kharif season Pointed Gourd: Epila									hna beetle insect						Duration: 100-120 days								
Weather warning	Months			М	arch			Α	pril		May						June					July		
	Max. Temp. (°C)					Hi	igh temper	ature (35	°C); 25°0	C temperature	favored both fe	cundity and	d food con	sumption	of the Epil	achna bee	tle as well as larval dev	elopmer	nt					
	Min. Temp. (°C)									P	opulation decrea	ased with th	ne tempera	ature fall b	elow 23 o									
	RH Max (%)									Hig	h relative humidi	ty favored	Epilachna	beetle (7	0% or abo	ve)								
	RH Min (%)												0%											
	Rainfall (mm)						Rainfall is favorable for the activity of Epilachna beetle pop								tle popula	ulation								
	Std. Week	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	
lar	Rainfall (mm)	5.0	-	4.0	-	8.0	8.0	12.0	26.0	28.0	37.0	42.0	45.0	58.0	59.0	74.0	70.0	92.0	76.0	93.0	80.0	91.0	76.0	
norm:	Max. Temp. (°C)		-	_	-	32.9	33.9	34.3	34.0	33.6	33.1	33.6	33.4	33.3	33.9	33.4	32.7	32.3	32.3	32.0	31.9	32.0	32.1	
Weekly r weath	Min. Temp. (°C)					21.1	22.3	23.0	23.0	23.1	23.3	23.9	24.3	24.6	25.0	25.9	26.0	26.1	26.1	26.0	26.1	26.0		
			89.1		_	87.7	88.0	88.3	90.6	90.9	92.4	92.4	92.7	93.0	93.7	94.1	95.1	95.1	95.6	95.9	95.9	95.9	95.9	
	RH Min (%)		36.9		_	45.3	44.4	45.9	50.7	54.9	57.3	58.7	60.0	63.4	64.6	66.7	70.4	72.4	73.9	73.3	74.1	74.4	73.3	
	SS hr (hrs)	60.0	59.0	61	58.0	56.0	56.0	56.0	55.0	52.0	48.0	50.0	49.0	46.0	44.0	39.0	33.0	28.0	26.0	28.0	31.0	29.0	28.0	
G	Spo	outing	& See	edling		vegetative growth					Flowering & fruiting						Harvesting							
Ins	sect Life cycle					(2 to 5 days) (4 to 9 days) (9 to 15 days)				Adults (3 to 10 days)	Eggs (2 to 5 days) (4 to 9 days) Pupa (9 to 15 days) (3 to 10 to													