



NATIONAL AGROMETEOROLOGICAL ADVISORY SERVICE BULLETIN

25 September 2019 (WEDNESDAY)
[FOR THE PERIOD 25.09.2019 - 29.09.2019]



DISCLAIMER

Under the Agro-meteorological Information System Development Project, Agromet Advisory Service Bulletins at National & Sixty Four (64) districts have been prepared on experimental basis. These bulletins are being displayed through the website of Department of Agricultural Extension for valuable comments/ suggestions from all concerned.

Please Contact: Farhana Hoque, Sabuj Roy | E-mail: pdamisdo@dae.gov.bd
Phone No.: 02-55'028414, 02-55028418

ISSUED BY

Agro-Meteorological Information Systems
Development Project
Department of Agricultural Extension (DAE)
Ministry of Agriculture
In collaboration with Bangladesh
Meteorological Department

Salient Agromet Advisories

As per the weather forecast received from the Bangladesh Meteorological Department (BMD), the axis of monsoon trough runs through Madhya Pradesh, Odisha to Assam across Gangetic West Bengal and southern part of Bangladesh. One of its associated troughs extends up to North Bay. Monsoon is fairly active over Bangladesh and weak to moderate over North Bay. During next 24 hours light to moderate rain/thunder showers accompanied by temporary gusty wind is likely to occur at many places over Rangpur & Khulna divisions and at a few places over Barishal, Dhaka, Rajshahi, Mymensingh, Sylhet & Chattogram divisions with moderately heavy to heavy falls at places over the country. As per the medium range quantitative district level weather forecast issued by BMD, a number of districts like Rangpur, Thakurgaon, Shariatpur, Panchagarh, Noakhali, Naogaon, Nilphamari, Narsingdi, Madaripur, Laksmipur, Joypurhat, Gaibandha, Dinajpur, Chapai Nawabganj, Bhola & Barishal is likely to receive relatively more rainfall (more than 50mm) during next five days compared to other districts in the country. As a result, it is advised not to apply irrigation, pesticides and fertilisers and also not to carry out intercultural operation in these districts where relatively good amount of rainfall is expected during next few days. Day temperature may rise slightly and night temperature may remain nearly unchanged over the country. As per the information received from Bangladesh Water Development Board (BWDB). All the major rivers are in rising trend except the Brahmaputra-Jumana. The Brahmaputra-Jumana rivers may continue falling while the Ganges-Padma rivers may continue rising in next 24 hours. The Surma-Kushiyara rivers may rise in next 24 hours. Considering the realized weather along with weather forecast, following salient agromet advisories are issued for different districts, where good amount of rainfall is likely for next five days, in the country on the standing crops, livestock, poultry and fishery.

Aus Paddy

- Drain out water completely 15 days before harvesting
- Harvest the crop after the current spell of rainfall.

Aman Paddy

- Since there is possibility of getting good amount rainfall in the coming 5 days, it is advised to repair the bunds in the paddy fields to conserve rain water.
- Maintain 5-7 cm standing water up to maximum tillering stage utilizing the good amount of rainfall received during last four days and expected more rainfall for next five days. At panicle initiation stage, 2-5 cm water level should be maintained.
- All the intercultural operations should be done after the current spell of rainfall
- First weeding should be done 10-15 days after transplanting. Second weeding should be done 30-35 days after transplanting. Weeding may be done either by hand or using weedicides. 2-4 D Amine or Butachlor may be used for weeding.
- Apply 1/3rd nitrogen as top dressing 15-20 days after transplanting. Last 1/3 nitrogen as top dressing should be applied 5-7 days before panicle initiation. All these operations should be done after the current spell of rainfall.
- Continuous monitoring of the economically harmful insects like stem borer, rice hispa, rice caseworm, gall midge should be done. For this light trap should be used. Light trap should be placed at some distance from the crop and bulb should be lightened from evening 06.30 PM to 10.30 PM and collected insects should be destroyed in the morning hours.
- There is chance of attack of leaf folder /rice hispa in seedling and tillering stage. If there is one damage leaf per hill due to attack of leaf folder or appearance of one adult hispa per hill is observed,

the farmers are advised to spray Chloropyriphos 20EC or Monocrotophos 40EC @ 1.5ml per litre of water.

- In some places, crops are attacked either by stem borer or leaf eating insects. To control broadcast granular insecticides like Carbofuran 3G (at 12 kg per acre) or Phorate 10G (at 4 kg per acre).
- Regularly monitor the rice fields for plant hopper infestation. If 5 or more plant hoppers per hill are observed, spray Chess 50 WG @ 170 g/acre or Confidor 200 SL @ 40 ml/acre.
- Due to persistence of high humidity and cloudy weather. false smut may appear in rice, spray with Kocide 46 DF (Chopper hydroxide) @ 500 g in 200 litres of water per acre to control false smut and after 10 days of its application spray Tilt 200 ml in 200 litres of water.
- All plant protection measures should be done after the current spell of rainfall.
- In low land areas, still there is scope for transplanting the crop after receding water. As there is delay in transplanting, cultivation of BR22, BR23, BRR1 DHAN 38, BRR1 46, Binashail, Nigershal and local varieties may be suitable under such circumstances.

Vegetables

- Provide adequate drainage facilities to drain out excess water from the field.
- Provide staking of vegetables to prevent the crops from lodging due to strong winds.
- All plant protection measures should be done after the current spell of rainfall

Ladies finger At fruiting stage there is a chance of incidences of leaf spot, Jassid, white fly In case of spots on leaves with grey and dark margin apply carbendazim 50% WP @ 1 g/lit of water. For managing jassids and white fly Thiomithoxam 1gm/3lit. of water. Cleanliness to be maintained in the field.

Cucurbits : At fruiting stage, there is a chance of incidences of fruit fly, white fly, mosaic Collection and destruction of fruit fly infested fruits. Installation of pheromone trap @ 2/Bigha. For white fly apply clothianidin 50 WDG@ 3g/10lit. of water. For mosaic uproot the infected plant and spray systemic insecticide. Due to poor pollination fruit setting in such case apply Boron 20% @ 2g/lit. of water to enhance fertilization at fruit setting. Do not use any pesticide and insecticide during flowering and fruit setting.

Brinjal: At vegetative & fruiting stages, there is chance of fruit and shoot borer, nematode, white fly, Jassid, fruit rot. For the management of fruit and shoot borer one application of coragen @ 4 ml/tank. Cleanliness to be maintained. Apply Thiomethoxam 0.3 SG @1g/3lit. of water to manage sucking pests infestation. For nematode problem apply Fipronil 0.3G @1.5 Kg/Bigha followed by a light irrigation or Nimitz 480 EC@ 4-8 lit/ha. could be applied against root knot nematode. For white fly and jassid, apply Acetamiprid 20% SP@1g/3lit or Thiomithoxam 0.3 SG 1g/3lit of water. Avoid touching of fruits with soil to avoid rotting of fruits.

Bitter gourd & Pointed gourd: At vegetative growth and under the present weather condition, attack of Downy mildew may be seen in the field. Mancozeb or Chlorothalonil 2 g/lit twice at 10 days interval may be sprayed to manage the attack.

Horticultural crops

- Planting of new saplings of fruit plants such as mango, litchi, jackfruit, guava, blackberry, custard apple, lemon is advised. If damaged by heavy rainfall, fill the gaps by replanting.
- The farmers are advised to take up plant protection measures carefully in horticultural crops like, pomegranate for bacterial blight and lime for leaf minor.

- The current weather is congenial for fruit blight and fruit rot disease in pomegranate, to control, spray Mancozeb 600g and Carbendazim 100 g @ 200 litre of water. . To control thrips in pomegranate, spray Spinosad 2.5 SC @ 10 ml per 10 liters of water.
- Use fruit fly trap to control fruit fly damage in guava orchards and cucurbits. Farmers are advised to remove foot rot infected plants from rice field and avoid ponding of water to check further spreading of infection.
- To control fruit blight and fruit rot disease in pomegranate, spray Mancozeb 600g and Carbendazim 100 g @ 200 litre of water.
- All plant protection measures should be done after the current spell of rainfall

Coconut

- Prepare pit for plantation of saplings of coconut
- Due to prolonged rains, there is a chance of bud rot in coconut. Keep three perforated Mancozeb sachets (5 g per packet) around spindle leaves as a prophylactic measure against bud rot
- Collect all the fallen coconuts from the field to avoid sprouting.

Banana

- Provide mechanical support to young banana to prevent the crops from lodging due to strong winds
- Rake up the accumulated mud from the root zone of banana for better aeration.
- Plantation of banana crop should be done
- Due to monsoon showers, there is a chance of Sigatoka disease in banana. Cut and burn all severely affected leaves. Spray 1% Bordeaux mixture with sticker soon after the appearance of symptoms (5 to 6 sprays at fortnightly intervals). Spray 1% petroleum based mineral oils. Spray 0.1% carbendazim or 0.2% Mancozeb alternatively @ fortnightly intervals. Spray Azoxystrobin, Tebuconazole, Difenconazole (0.1%)
- There is a chance of erwinia rot in banana due to heavy rainfall in this season. As a precaution, apply lime in the banana basin or bleaching powder in the irrigation channels. To enhance the disease resistance, drench the basin with Pseudomonas @ 20g per litre of water. Affected plants can be protected by drenching the basin with 5 g bleaching powder or 3 g Copper oxychloride or 2g Copper hydroxide per one litre of water.
- All plant protection measures should be done after the current spell of rainfall.
- Severely affected plants should be cut and removed to prevent the spread of disease.

Sugarcane:

- Need based intercultural operations and earthing should be done.
- Irrigation can be given at weekly interval (for medium type of soil). In heavy soils, 10-15 days interval.
- If water scarcity is there irrigate the crop in alternate row, to save the water. Drip irrigation can save the water up to 40 per cent.
- Application of sulphur in the form of Gypsum @ 500 kg /ha to sulphur deficient soils to increase the cane yield and juice quality.
- Proper drainage system in the sugarcane fields should be maintained to minimize outbreaks of Pycnia pests

- There is a chance for early shoot borer incidence in sugarcane, farmers are advised to spray Chlorpyrifos@2.5ml/L or Monochrotophos@1.6ml/L for control.
- To control red rot disease in sugarcane, avoid water stagnation in the field and removing of the affected plants should be practiced.
- To minimize infestation of top shoot borer in sugarcane, spray Lambda cyhalothrin @2ml/litre of water.
- All plant protection measures should be done after the current spell of rainfall.

Turmeric:

- High relative humidity may cause leaf spot in turmeric, spray Mancozeb 1 kg/ha after the current spell of rainfall.
- To control Rhizome rot disease turmeric, drench the affected area with 0.3% Mencozeb 75 WP @ 2g/l of water after the current spell of rainfall.

Betelvine:

- Strengthen the fencing of boundary of betel vine so that the boundary cannot be broken due to strong wind and thunderstorm.
- Keep the drainage channels clean and arrange proper aeration in the betel vine gardens.
- No spraying of copper fungicide for control of foot and root rot should be given until all the water will removed.
- Pluck muddy leaves from the lower parts of betel nut plants.
- Farmers can continue to plant the cuttings of betel vine in the main field. Select disease free cuttings and before planting treat them with Bordeaux mixture 0.5% and Streptocycline 500 ppm for half an hour. Before planting, drenching of soil with Mancozeb 75 WP (2.5 gram per litre of water) is advocated after the current spell of rainfall

Cotton: Sowing stage

- Continue sowing of cotton.
- Under humid weather conditions, there may be increase in population of insect pest of cotton. Farmers are advised to install pest specific pheromone traps in the fields for monitoring the insect. If sucking pest's population is above ETL, spraying of Imidacloprid 17.8 % SC @ 1.0 ml / 3 litre or Trizophos 40 EC @ 1.0 ml/litre of water is advised against all type of borers after the current spell of rainfall.
- Due to variable weather conditions, high humidity and favourable temperature for the chances or infestation of white fly. Farmers are advised to keep vigil on leafhopper, thrips white fly infestation. If infestation of white fly above to the economical threshold level (ETL) than 400 ml dimethoate (rogor) 30 E.C. or 500 ml oxydemeton-methyl 25 E.C. (metasystox) spray of 1 liter Nimbecidine mixed with 250 liter water per acre, when sky remain clear. For control of leafhopper infestation the spray 40 ml imidacloprid (Confidor) 200SL mixed with 120-150 liter water per acre, during clear weather

Ginger:

- Allow proper drainage from raised beds of ginger with necessary arrangements. Apply hexaconazole @1ml/lit of water as foliar spray if ash colour spots develop on leaves. Spray with (Melanayl 8%+mancozeb 4) @ 2g/lit of water as soil drenching followed by caoxychloride @ 4g/lit of water after the current spell of rainfall.

- Weather is congenial for soft rot in ginger, drench the soil with 1% Bordeaux mixture/0.3% Mancozeb and apply green leaf mulching to prevent soil erosion and to increase the organic matter content in the soil
- Under excessive moisture, there is every possibility of rhizome rot in ginger and turmeric. To manage the disease, provide adequate drainage and apply Trichoderma Harzianum @ 50 g/bed along with FYM/ neem cake and undertake mulching.

Livestock, Poultry & Fishery

- Hygiene should be maintained around cattle shed. Avoid water logging in cattle shed.
- Vaccinate cattle and buffalos against the rainy season diseases. Protect cattle from sun.
- Vaccinate the livestock for Black Quarter and foot and mouth diseases. Sheep/goats must be dewormed for maximum weight gain and good health. There is chance of Raniket, gumbra diseases in poultry; provide antibiotic drugs through water or through feed.
- During the monsoon season, to prevent the livestock from worm infestations, it is advised to give Piperzine or Mebendazole at recommended dose after proper consultation with veterinary doctor.
- During rainy days, goats may be affected with viral enteritis. As and when diarrhoea is noticed, immediate treatment should be done at the veterinary dispensary and do not allow the goats to graze outside.
- Protect poultry houses from extreme rain/wind by covering with polythene sheets on the sheds. Provide vitamin B complex @ 7ml/litre in drinking water daily in broiler for healthy growth. Litter should be change periodically and provide clean drinking water and changed the water every day.
- To avoid insect infestation in nursery ponds, apply soap oil emulsion (mixture of cheap soap @ 2.5 kg and kerosene oil @ 13 litre per bigha of pond) and apply gently in nursery ponds.

Detail District-wise Agromet Advisories are available in BAMIS Portal (www. <https://www.bamis.gov.bd/>)

Realized Weather at different locations in the country

Rainfall for last 24 hours (till 06 AM on 25 September'19) and maximum temperature on 24 September, minimum temperature on 25 September'19 is given below:

Name of Divisions	Name of Stations	Rain fall (mm)	Max. Temp (°C)	Min. Temp (°C)	Name of Divisions	Name of Stations	Rain fall (mm)	Max. Temp (°C)	Min. Temp (°C)
Dhaka	Dhaka	08	31.6	24.8	Rajshahi	Rajshahi	21	33.8	25.4
	Tangail	01	32.5	24.5		Ishurdi	10	32.5	25.2
	Faridpur	05	31.8	23.7		Bogura	13	30.8	25.0
	Madaripur	02	31.2	24.6		Badalgachhi	04	30.6	24.4
	Gopalganj	09	30.3	24.5		Tarash	02	31.7	25.0
	Nikli	25	33.0	24.3					
Mymensingh	Mymensingh	16	31.3	24.3	Rangpur	Rangpur	160	27.2	24.4
	Netrokona	37	29.6	23.5		Dinajpur	44	30.3	24.0
						Sayedpur	07	28.2	24.1
Chattogram	Chattogram	00	33.1	25.3		Tetulia	23	28.2	23.8
	Sandwip	Trace	33.2	25.4		Dimla	17	29.0	23.3
	Sitakunda	00	33.3	26.5		Rajarhat	10	26.8	24.0
	Rangamati	02	33.5	24.4	Khulna	Khulna	02	32.0	25.0
	Cumilla	16	32.0	23.5		Mongla	23	30.0	24.6
	Chandpur	43	32.2	24.0		Satkhira	26	34.8	24.0
	M.Court	19	31.5	24.0		Jashore	08	32.6	25.0
	Feni	03	32.5	24.4		Chuadanga	04	32.0	24.7
	Hatiya	25	31.2	24.3		Kumarkhali	08	xx	25.2
	Cox's Bazar	00	33.0	26.3	Barishal	Barishal	18	31.2	24.8
Kutubdia	01	32.8	25.3	Patuakhali		16	30.6	24.5	
Teknaf	00	34.1	26.5	Khepupara		37	28.6	24.4	
				Bhola		61	30.9	24.6	
Sylhet	Sylhet	08	30.3	23.5					
	Srimangal	12	31.4	23.2					

Highlights:


- Country average bright sunshine hour was 6.33 hours per day during last week.
- Country average free water loss during the previous week was 3.65 mm per day.





Weather Forecast Valid for 24 Hours commencing 06 PM Today:





Forecast: Light to moderate rain/thunder showers accompanied by temporary gusty wind is likely to occur at many places over Rangpur, Rajshahi, Dhaka, Mymensingh, Khulna, Barishal & Sylhet divisions and at a few places over Chattogram division with moderately heavy to heavy falls at places over the country.

Temperature: Day and night temperature may remain nearly unchanged over the country.

Spatial Distribution of weather parameters for the Week ending 24 September 2019








Weather Forecast and Advisory for the Period of 22.09.2019 to 30.09.2019





Bright sunshine hour is likely to be from 4.00 to 5.00 hours per day during this week. Average free water loss during the next week is likely to be from 2.50 to 3.50 mm per day.

Light (04-10 mm/day) to moderate (11-22 mm/day) rain/thunder shower is likely to occur at many places over Dhaka, Rajshahi, Khulna, Barishal, Chattogram, Rangpur, Sylhet, And Mymensingh divisions, with chances of moderately heavy (23-43 mm/day) to heavy (44-88 mm/day) falls at places over the country during this period.

Day and night temperature may fall slightly over the country during this period.


District Wise Quantitative Weather Forecast for next Five days from 25 Sept. -29 September'19






Different Satellite Products over Bangladesh


NOAA/VIIRS BLENDED NDVI composite for the week number No. 37 (7 September -13 September) over Agricultural regions of Bangladesh




NOAA/ AVHRR BLENDED VCI composite for the week number No. 37 (7 September -13 September) over Agricultural regions of Bangladesh



NOAA/ AVHRR BLENDED TCI composite for the week number No. 37 (7 September -13 September) over Agricultural regions of Bangladesh




NOAA/ AVHRR BLENDED VHI composite for the week number No. 37 (7 September -13 September) over Agricultural regions of Bangladesh



Monitoring Meteorological Drought in Bangladesh using Standardized Precipitation Index (SPI)

It has been observed that during last four weeks as well as in August to September, mild wet conditions prevails in districts located in northwest as well as southeast parts of Bangladesh whereas the districts in south, northeast, west and central parts of Bangladesh are under mild to moderate dry condition.



Data source: Bangladesh Meteorological Department

RAINFALL AND RIVER SITUATION SUMMARY AS ON SEPTEMBER 25, 2019

(Source: Bangladesh Water Development Board)

Outlook:

- All the major rivers are in rising trend except the Brahmaputra-Jumana.
- The Brahmaputra-Jumana rivers may continue falling while the Ganges-Padma rivers may continue rising in next 24 hours.
- The Surma-Kushiyara rivers may rise in next 24 hours.

General River Condition

Monitoring Water Level Stations	93	Water Level Steady in last 24 hours	03
Water Level Rise in last 24 hours	66	Total not Reported	00
Water Level Fall in last 24 hours	24	Above Danger Level	00