

Agromet Advisory Bulletin for Bandarban District

		
		
<p>Agro-Meteorological Information Systems Development Project Component-C of BWCSR Department of Agricultural Extension</p>		
Date: 25th March 2020 Bulletin No. 131	Agromet Advisory Bulletin for Bandarban District (25th March to 29th March 2020)	

Weather Conditions for last four days (21st March to 24th March 2020)

Weather Parameters	21.03.20	22.03.20	23.03.20	24.03.20	Range
Rainfall (mm)	0.0	0.0	0.0	0.0	0.0-0.0 (0.0)
Maximum Temperature (° C)	30.6	30.4	30.4	29.8	29.8-30.6
Minimum Temperature (° C)	22.5	22.2	19.8	21.3	19.8-22.5
Relative Humidity (%)	61.0-93.0	36.0-85.0	50.0-91.0	57.0-96.0	36-96
Wind Speed (km/h)	7.4	5.6	5.6	9.2	5.55-9.25
Cloud Amount (Okta)	5	4	1	3	1-5
Wind Direction	West/Northwesterly	West/Northwesterly	West/Northwesterly	West/Northwesterly	West/Northwesterly

Weather forecast as per Bangladesh Meteorological Department for the next 5 days (25th March to 29th March 2020)

Weather Parameters	Range
Rainfall (mm)	0.0-0.0 (0.0)
Maximum Temperature (° C)	32.1-36.9
Minimum Temperature (° C)	16.4-17.9
Relative Humidity (%)	41.0-80.0
Wind Speed (Km/h)	2.3-3.1
Cloud Amount (Okta)	Partly Cloudy Sky
Wind Direction	West/Northwesterly

Agromet Advisories

Salient Weather Conditions & Agromet Advisories

Dry weather condition with partly cloudy sky will prevail. Day & night temperature is likely to increase slightly. As per the outlook issued by BMD for next 72 hours day temperature is likely to increase further. Detail Agromet advisories are given below.

Boro Paddy

Tillering to Booting stage

- Alternate wetting and drying maintain water level 5-7 cm up to the appearance of panicle initiation stage. Maintain 2-5 cm water level in booting stages.
- There is chance of stem borer in rice in the main field. If noticed, the pest may be identified and controlled by using light traps and also constant monitoring is required for its level of incidences. To attract the butterflies, place 5 Pheromone Traps/acre, collect the adults and destroy them by burying under soil or by burning. Also spray Carbofuran @10kg per ha
- Before spray the insecticide, drain out the irrigated water from the plots and
- Increased difference between daily maximum and minimum temperature (diurnal temperature) may favour the infestation of blast in main field as well as in the nursery. At peak tillering stage of the crop there is more chance of this fungal disease. Spray Nativo 75wg/ Trooper@ 0.6g/litre of water or Amister top 325 sp@ 1ml/litre of water
- For brown spot attack spray apply fertilizer management and spray Thiovit+Potash
- Spraying should be done during afternoon hours or at morning hours from 10:00 a.m. to 11:00a.m

Groundnut

- Monitor the incidences of pests (hairy caterpillar, jassid, thrips, leaf miner, spodoptera & termite) & diseases (leaf spot/tikka disease, leaf rust, foot rot/stem rot, collar rot, bud necrosis) and if noticed, take appropriate preventive measure.
- Current moderate temperature favours thrips infestation. Spray 400 ml Dimethoate or 50 ml Imidachlopid in 200 litre water per acre.
- Present weather conditions are favourable for leaf miner incidence in early sown groundnut. For control of leaf miner to spray Chloripyriophos 2.5 ml/l or quinalphos 2 ml/l of water sowing and
- Present weather conditions are favourable for spodoptera, jassid in groundnut. To control leaf minor and spodoptera, spray Chloripyriophos @ 2.5 ml (or) Quinalphos @ 2 ml or Thiodicarb 1 g per litre of water; for sucking pest, spray Monochrotophos @1.6 ml (or) Imidaclopid @ 0.3 ml (or) Dimethoate @ 2 ml per litre of water and for tikka leaf spot, spray Mancozeb @ 400 g + Carbendazim @ 200 g (or) Hexaconazole @ 400 g per acre.
- There is a chance of infestation of collar rot disease in early sown rabi groundnut crop. Spray of Thiophanatemethyl 1.5 g/liter water is advised.

Vegetables

- Prevailing weather conditions are favorable for thrips & aphids in chilli. Spray Emamectin benzoate 5% SG 4 g/10 litre or Dimethoate 30 % EC 1.0 ml/litre for the control of thrips & aphids.
- Intercultural operation is advised in early sown onion/garlic crop. Provide light and frequent irrigation at an interval of 10-15 days as per requirement.
- For the management of stem and fruit borer in brinjal crop, it is advised to install 10 Pheromone trap in one acre
- Avoid moisture stress conditions to the crops which are at flowering to fruiting stage.
- Considering the increase in temperature, apply light irrigation in standing crops and vegetables as per requirement.
- There is chance of blight disease in tomato and brinjal which are at fruiting stage. Apply Streptocyclene@3g/10 liter of water.

Horticultural crops

- Due to continuous dry spell, there are chances of termite attack in horticulture, forestry tree and shrubs. Hence, to control apply Aldrin termiticides for termites.
- Farmers are advised to keep vigilance on insect-pest and diseases and follow proper weed control measures and cover fruit seedling with rice straw (parali)/thatches in nursery.
- To avoid fruit fall in mango crop, it is advised to spray Planofix regulator 20 ppm and 1% Calbor micronutrient.
- Wherever flowering is delayed in mango, farmers are advised to give irrigation and spray potassium nitrate 10 g and Urea 5 g per litre of water to enhance the flowering.
- Avoid moisture stress conditions to the crops which are at flowering to fruiting stage.
- Apply coir compost to conserve moisture in banana
- Young banana plants of 3-4 months' age are prone to attack of pseudostem weevil; spray Chlorpyrifos 2 ml/litres of water.
- There is a chance powdery mildew disease incidence before flowering and after fruit formation in mango. To control, spraying of Carbaryl, 50WP @4g/litre of water or Imidachlorpid @ 0.3ml/ litre of water is advised.
- To protect mango plants from mealy bug, spray Imidaclopid 17.8 SL @ 1 ml /3 litres of water
- Mango is in tender fruiting stage and there is chance of hopper attack.If noticed, apply a mixture of Diethane M-45@2.5g/liter of water and Dimethoate@1.5ml/liter of water.
- Coconut harvested during November can be sown in nursery bed for raising seedling. Nuts should be placed horizontally with widest of the three segments upward which helps in germination and healthy growth of seedlings. In 1m wide nursery beds, the nuts should be placed at a spacing of 30cm x 30cm.

Jute:

- Land is to be ploughed and laddered 5-6 times to obtain a fine seed bed. In termite and cricket infested fields, the soil is to be treated with malathion 5% dust @ 30 kg/ha.
- If there is enough soil moisture, start sowing of seeds.

Livestock

- Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.
- Keep the animal shed clean and dry.
- Green fodder mixed with dry straw should be provided for feed.
- In case of skin disease, dairy farmers are advised to mix zinc oxide and vaseline in equal proportion and apply on affected parts.
- Keep away mosquitos from animal shed by suitable technique.
- At this time goat may be attack by Blister disease which is generally a viral disease that affect any age group of goat. The disease may be seen particularly in the facial/ nasal area. To protect from the disease wash with Potassium per manganate (PP) and consult nearby Veterinary Dispensary for anti-biotic cream and further treatment.
- Feed mineral especially calcium and vitamin supplement to the dairy cattle to have good productive and reproductive performance

Poultry

- Suggested to go for routine vaccination for poultry against the dreaded diseases /virus Vaccination (RDF1) of the one week old chicks against Raniket disease, darkplaigue and against Gamboro disease in two weeks old chicks may be carried out after consultation with local veterinarian.
- It is advised to align the long axis of poultry shed in east-west direction. This will prevent the direct sunshine over the birds.
- Regularly inspect the droppings of the poultry bird for any disease symptoms and immediately separate/remove the diseased and dead birds from the healthy ones.

Fishery

- Because of increase in temperature, it is advised to add cow dung and lime or rice bran and mustard cake in the pond as a natural food for fishes @ 5 kg/day/ acre area of pond.
- Maintain 1 to 1.5mt average depth of water in the pond.
- As the temperature is rising, farmers can start feeding their fishes with available fish feed or they can feed their fishes with a mixture Rice bran and MOC in 1:1 ratio @3% of body weight of the total biomass in their ponds.
- To avoid bacterial disease liming should be done at appropriate dose.