




Agromet Advisory Bulletin for Bandarban District

		
		
<p>Agro-Meteorological Information Systems Development Project Component-C of BWCSR Department of Agricultural Extension</p>		
Date: 29th July 2020 Bulletin No. 167	Agromet Advisory Bulletin for Bandarban District (29th July to 02nd August 2020)	

Weather Conditions for last four days (25th July to 28th July 2020)

Weather Parameters	25.07.20	26.07.20	27.07.20	28.07.20	Range
Rainfall (mm)	0.0	0.0	0.0	35.0	0.0-35.0 (35.0)
Maximum Temperature (° C)	33.1	33.0	33.0	33.3	33.0-33.3
Minimum Temperature (° C)	26.2	26.1	27.8	27.0	26.1-27.8
Relative Humidity (%)	63.0-90.0	67.0-98.0	67.0-92.0	66.0-96.0	63-98
Wind Speed (km/h)	13.0	11.1	7.4	14.8	7.4-14.8
Cloud Amount (Okta)	6	6	6	7	6-7
Wind Direction	South/South-westerly	South/South-westerly	South/South-westerly	South/South-westerly	South/South-westerly

Weather forecast as per Bangladesh Meteorological Department for the next 5 days (29th July to 02nd August 2020)

Weather Parameters	Range
Rainfall (mm)	0.7-12.7 (27.7)
Maximum Temperature (° C)	30.5-31.4
Minimum Temperature (° C)	23.5-24.4
Relative Humidity (%)	85.0-96.0
Wind Speed (Km/h)	1.5-2.5
Cloud status	Partly Cloudy Sky
Wind Direction	South/South-westerly

Agromet Advisories

Special Agromet Advisories for CORONA virus (COVID-19) transmission in Bangladesh

Farmers and all other farm managements are advised to maintain the social distancing (at least 3 meters), cover their faces with masks and follow other guidelines of Government of Bangladesh to avoid infection and social transmission of CORONA virus (COVID-19).

Salient Weather Conditions & Forecast

The axis of monsoon trough runs through Punjab, Haryana, Uttar Pradesh, Bihar, Sub-Himalayan West Bengal to Assam across Northern part of Bangladesh. One of its associated troughs extends up to North Bay. Monsoon is fairly active over Bangladesh and weak to moderate over North Bay. As per the weather forecast received from the Bangladesh Meteorological Department (BMD) during next 24 hours, light to moderate rain/thunder showers accompanied by temporary gusty wind is likely to occur at many places over the district with moderately heavy to heavy falls at places over the district. Night temperature may remain nearly unchanged over the country. As per the outlook issued by BMD for next 72 hours, little change is likely. During last four days, no rainfall occurred and as per the quantitative medium range weather forecast, trace to moderate rainfall is likely during next five days. Detail Agromet advisories are given below.

Aus Paddy

Tillering to Booting stages

- Maintain in 3-4 inches water level. Maintain 5-7 cm water level from panicle initiation stage.
- To control infestation of leaf blast & leaf spot in Aus rice, spray Carbendazim @2g/litre of water.
- Due to high humidity (warm and humid weather) possibility of pest and disease build up is more in crop. Take prophylactic measures accordingly.
- To control of insects like stem borer gall midge and white & brown hopper apply Corbofuron 3 G@33 kg or phorat 10G @ 10 kg per /hectare and for cut worm Clorpyrifos or Daiclorovash.
- Since cloudy weather is prevailed, there is a chance of leaf folder attack in Paddy. It can be controlled by using Trichogramma cards.
- To save the rice crop from sheath blight, keep the bunds of the field clean by removing grasses.

Aman Paddy

- If the rice seedbed is underwater, it is expected to be damaged and advised for preparation seedbed in trays (if possible) in some raised structure. The farmers who had already completed their transplanting are advised to maintain drainage.
- Complete transplanting of rice seedlings at the earliest as sufficient water is there in the field; maintain proper drainage in field/nursery to avoid flooding due to continuous rain.
- Ensure weed free seed bed.
- Farmers may start the preparation of main land for transplanting after getting sufficient rainfall.

- At the last stage of land preparation of field for transplanting of Aman paddy, apply per hectare 90 kg TSP, 70 kg MOP, 11 kg Zinc, and 60 kg gypsum.
- Due to substantial decrease in sunshine hours with occasional rain there is chance of attack of Yellow stem borer in Aman rice; to manage the attack, spray Carbofuran @10kg per ha.
- Transplanting of 25-30 days old nursery in the main after getting sufficient rainfall.
- Arrange the bunding tightly so that no drainage can take place after light rain or moderate rain
- Transplanting of aman paddy in the main field is going on due to sufficient amount of water present in the field. Thereby, it is recommended that before transplanting the seedling in the main field, farmers are advised to uprooted seedlings are washed and then the root portion is dipped in 0.02% solution of chlorpyrifos (1ml/lit of water) along with 1% urea (10gm/lit of water) for three hours as protective measure against stem borer, gall midge and hopper
- Avoid deep planting for better crop establishment and fill up the gaps within a week. Maintain 5-7 cm standing water up to maximum tillering stage. Keep the rice nursery and field bunds free from weeds either by hand weeding or by application of herbicide. Apply recommended herbicide as per recommendation within 1-3 days of transplanting.
- Apply 1/3rd nitrogen as top dressing 15-20 days after transplanting and after the current spells of heavy rainfall.
- After reduction of rainfall activity mechanical weeding can be started to control the weed at initial stage. For chemical control (only if the sky is clear and no rainfall 7 to 8 hours after application) you can spray Pretilachlor 50% EC @500 gm per acre within 2-5 days after transplanting. Due to stagnation of water in the field and high relative humidity Sheath blight - Fungal disease is common. Excess Nitrogen fertilizer application can accelerate this disease. August to October is the high time for this disease attack. Control: 1) Drain out excess water from infected field. 2) Reduce the dose of Nitrogen fertilizer. 3) Bio fungicide like Trichoderma, Pseudomonas Fluorescens is also good. 4) Spray Propiconazol+ Difenconazole 1ml/liter of water or Carbendazim + Flusilazole 1.5 ml/lit or Carbendazim (Bavistin) 1 gram/liter of water
- Under congenial condition, following aman variety may be sown.
- BRRRI DHAN 30, BRRRI DHAN 32, BRRRI DHAN 39, BRRRI DHAN 49, BRRRI DHAN 62 BRRRI DHAN 71, BRRRI DHAN 72, BRRRI DHAN 75, BRRRI DHAN 80, BRRRI DHAN 87, BRRRI DHAN 90, BRRRI DHAN 93, BRRRI DHAN 94, BRRRI DHAN 95, BINA DHAN 11, BINA DHAN 16, BINA DHAN 22
- Due to substantial decrease in sunshine hours with occasional rain there is chance of attack of Yellow stem borer in Aman rice; to manage the attack, spray Carbofuran @10kg per ha.

Vegetables

- Due to continuous heavy rainfall and water stagnation the vegetable crops like Brinjal, Pointed gourd, Bottle gourd, Ridge gourd, Chili, etc suffer from rotting. Make proper drainage channels to drain out water. Avoid spraying and application of fertilizer till the clear sky.
- Make arrangement to avoid plant damage (particularly staked creeper) from anticipated gusty winds. Make arrangement to raise new orchard during monsoon season.
- If seedlings of brinjal are ready for transplanting, transplant the seedlings on ridges and furrow with spacing of 60 x 60 cm. Sowing of kharif vegetables such as lady's finger, pumpkin, cucumber, sponge gourd, bottle gourd.

- In pumpkin, ridge-gourd, pointed gourd and cucumber, there is a chance of attack of Red-Pumpkin Beetle during current weather condition. To control, apply Dimecron or rogor (1ml per litre of water).
- The prevailing weather conditions are congenial for incidence sucking pests in vegetables. To control, spray Dimethoate @ 2 ml or Acephate @ 1.5 g per litre of water. Undertake nursery sowing of Tomato, Brinjal and Chillies and direct sowing of Gourds, Beans and Bhendi with assured irrigation.
- At flowering to fruiting stage of bitter gourd, there is chance of Rotting at the base of flower, apply SAF @2g/liter of water.
- Weed management should be done in brinjal, tomato, lady finger and cauliflower.
- Need based intercultural operations should be done in tomato, brinjal, okra and other vegetable crops.
- Undertake nursery sowing of Tomato, Brinjal and Chillies and direct sowing of Gourds, Beans and Bhendi.
- Collect Maskali and winter vegetable seed

Horticultural crops

- Farmers are advised to go for inter cultural operations in mango orchard.
- The farmers are advised to take up plant protection measures carefully in horticultural crops like, pomegranate for bacterial blight and lime for leaf minor.
- For the orchards of the guava, add 20-25 kg cow dung and 50 gms of heptachlor in the soil extracted from the pit and fill it again. For mango, amla and berry orchards add 30 kg of cow dung, 250 grams of single super phosphate and 50-100 g heptachlor in the soil extracted from the pit and fill it again. These operations may be done.
- Farmers are advised to start digging of pit of appropriate size and distance for new orchard establishment. Prepare pit for plantation of saplings of mango, guava and coconut as sufficient rainfall has been received.
- Plantation of banana crop should be done after getting sufficient amount of rainfall. Intercultural operation should be done in banana orchard.
- Provide mechanical support to banana, to prevent the crops from lodging due to strong winds.
- Due to deluge with rainy days farmers are advised not to apply fertilizer to the banana plant. Moreover, ensure to harvest the mature banana at proper stage.
- Due to monsoon showers, there is a chance of Sigatoka disease in banana. As a precaution, spray Pseudomonas 20g/litre. In case of severe attack, spray 2 ml hexaconazole or 1 ml propiconazole per one litre of water by mixing it with a sticker and spray on both the surface of the leaf.
- Attack of Papaya mealy bug on papaya fruit can be controlled by dusting of Chlorpyrifos 1.5% dust or Malathion 5% dust around the healthy plants; also destruct ant colonies with drenching of Chlorpyrifos 20 EC @ 2 ml/litre.

Jute:

- Harvesting and retting of jute may be done preferably at pre-flowering stage (120 days after sowing) to get better quality fibre. At present the water temperature in the retting tanks/ nalas may ideal for retting of jute. Early and timely sown capsularis jute may be harvested during this week and kept in the field in standing position for 3-4 days before retting for proper defoliation.
- If there is stagnant water in the jute field and there is no scope to drain out the water within 2-3 days, it is advised for immediate harvest the matured crop and put in water for retting.

Betel vine:

- There is chance of incidences of Phytophthora root rot and chital disease. For Phytophthora root rot apply 1% Bordeaux mixture at the base at 1-month interval. For Chitla apply 0.5% Bordeaux mixture at 15-20 days' interval.
- Maintain free air movement inside the garden.
- All the plant protection measure should be taken after the current spell of rainfall.

Livestock

- Do not allow animals to get wet during rainy season. Apply sufficient amount of Calcium Carbonate or bleaching powder to avoid germs. Never let a cow or goat graze in a place where there is standing water. It should be noted that cows or goats should not take paddy in any way at this time. If it takes a lot of paddy and the stomach starts to swell, then feed a spoonful of mustard oil. It will stop the production of gas in the stomach. In addition, blotocil or blotonil should be given in certain doses. If necessary, take the advice of a nearby veterinarian.
- It is very important to provide a particular amount of dewormers for both roundworm and tapeworm to any cattle, goats,
- Livestock may be affected by foot and mouth disease due to rainy weather. Following measures may be taken up.
- Feed them only dry food.
- Keep the shed clean and dry.
- If wound found at foot and mouth wash the wound with potassium per manganate solution.
- For control of biting fly, hang the alovera plant in the animal shed.
- Farmers who have not vaccinated their goat against Enterotoxaemia last month should vaccinate the goats strictly this month.
- If any malady symptom is shown in the animal then immediately consult with the veterinary doctor for proper treatments

Poultry

- Never allow chickens or ducks to eat wet or damp feed during this rainy season. It is very important to provide a particular amount of dewormers for both roundworm and tapeworm to poultry at this time.
- Apply lime bleaching powder and copper sulphet solution spray before placing them in farm
- Keep the chicks thin layer of litter
- Follow proper vaccination at proper interval
- Apply dry food and clean tap water for drinking purpose
- Prevailing hot and humid weather condition may lead to gastrointestinal parasites infection to the poultry. Clean sanitary conditions of poultry sheds are very much important

Fishery

- Apply the required quantity of calcium carbonate or lime in the submerged ponds. Check the overflow of water, fishes may drain out with overflowed water, for this Use fishing net in the possible exit points. Don't use excess feed to avoid water pollution.
- Repair damaged dykes of fisheries ponds and take care so that fishes do not escape. Maintain outlet pipe from the pond so that excess water pass away and the water level is maintaining.
- Fish grower framers are advised to collect the healthy fingerlings from the authentic sources as there is enough water in the fish pond.
- Remove the unwanted fishes and predator by repeated netting before leaving the new fingerling in the fish pond. This is time for release of new batches of fishes in fish ponds.
- Select healthy, diseases free breeds of fishes from authentic sources after consulting with fishery officials. Before releasing, the fingerlings should be treated with Potassium permanganate (@ 3-5%.