



NATIONAL AGROMETEOROLOGICAL ADVISORY SERVICE BULLETIN

25 August 2021 (WEDNESDAY)
[FOR THE PERIOD 25.08.2021 – 29.08.2021]



DISCLAIMER

Under the Agro-meteorological Information System Development Project, Agromet Advisory Service Bulletins at National & Sixty Four (64) districts have been prepared on experimental basis. These bulletins are being displayed through the website of Department of Agricultural Extension for valuable comments/ suggestions from all concerned.

Please Contact: Farhana Hoque, Sabuj Roy | E-mail: pdamisdp@dae.gov.bd
Phone No.: 02-55`028414, 02-55028418

ISSUED BY

Agro-Meteorological Information Systems
Development Project
Department of Agricultural Extension (DAE)
Ministry of Agriculture
In collaboration with Bangladesh
Meteorological Department

Special Agromet Advisories for CORONA virus (COVID-19) transmission in Bangladesh

Agromet Advisories

Farmers and all other farm managements are advised to follow the guidelines of Government of Bangladesh to avoid infection and social transmission of CORONA virus (COVID-19). Precautions and safety measures should be taken up to prevent the Carona virus spread. Simple measures include social distancing, maintaining personal hygiene by washing hands with soap, wearing of face mask, drink hot water, stay at home and cleaning of implements and machinery. Farmers should not work in a group; consult with a doctor in case of any symptom. Please stay at home does not visit the field until it is very necessary.

Salient Weather Conditions & Forecast

As per the weather forecast of Bangladesh Meteorological Department (BMD) during next 24 hours' light to moderate rain/thunder showers accompanied by temporary gusty wind is likely to occur at most places over Rangpur, Rajshahi, Mymensingh, Sylhet & Dhaka divisions and at many places over Chattogram, Khulna & Barishal divisions with moderately heavy to heavy falls at places over the country. Day and night temperature may fall slightly over the country. As per the outlook issued by BMD for next 72 hours' is likely little change. During last four days, light rainfall to very heavy rainfall occurred many of the district of the country and as per the quantitative medium range weather forecast; many of the districts of the country are likely to receive moderate rainfall to very heavy rainfall in the next five days. Detail Agromet advisory are given below:

Aus:

- Maintain the required water level in the land.
- Insecticides of carbosulfan group can be applied to control thrips, green leaf hopper and Rice Bug insects.
- Catch and kill the insects with hand nets to control the attack of Rice Hispa insects. Apply Carbaryl / Chlorpyrifos group pesticides if there are 4 adult Rice Hispa insects or 35% damaged leaves in each bunch.
- Apply pesticides after current spell of rainfall.
- Drain out the water from the rice field 15 days before harvest.
- Collect mature crops after rains and keep them in a safe place quickly.

Sugarcane

- Sugarcane stems can be attacked by red rot disease. Infected trees should be removed from the land along with bushes as soon as the disease is seen. Water should be removed from the sugarcane field very quickly.
- To control the Sugarcane Stem Borer, remove the soil from the base of the sugarcane with a spade. Drain out accumulated water from the land. Old dried leaves should be removed from the trees, collected and burnt or buried under the ground. Egg parasitic hornet Trichogramma Chilonis should be released in the affected land at the rate of one gram per hectare per week (approximately 50,000). Apply Virtako 40 WG (Thiamethoxam + Chlorantraniliprole) to the

infested sugarcane bush in the approved dose for control of stem borer insects with pesticides or Cartop granular pesticides such as Nokotap 6G cut the shallow drains on both sides of the sugarcane row and sprinkle in the drains then cover with soil or sprinkle at the base of the tree.

- Drain out excess water from the land.
- All the plant protection measure should be taken after the current spell of rainfall.

Vegetables

- In case of infestation of *eggplant* tip and fruit borer insects, cut and destroy the affected tip along with the insects. Insect reproduction is likely to be greatly reduced by using pheromone traps. Use spray of approval dose of insecticides, if necessary or recommended local organic insecticides may be used.
- Arrange for hand pollination of pumpkin vegetables. If beetles appear, kill them by hand in the morning and afternoon.
- In case of fly infestation in pumpkin, use pheromone and poison traps together. Pesticides of Alpha-Cypermethrin group can be used.
- In case of spider infestation, soak 1 kg half-broken neem seeds in 20 liters of water for 24 hours and spray the water (after chopping) on the underside of the leaves. If the infestation is severe, spray the spider mite Omit 57 EC at the rate of 2.0 ml per liter of water or Vertimek 1.8 EC at the rate of 12 ml per 10 liters of water.
- Drain out excess water from the land.
- Apply pesticides after current spell of rainfall.

Horticultural crops

- At this time it is suitable for planting seedlings. After planting the seedlings, tie the seedlings with hard poles.
- In case of symptoms of Sigatoka disease on banana leaves, 0.5 ml of Score or 2 g of Noin or Bavistin or 0.1 ml of Econazole / Folico should be mixed in every liter of water and sprayed 2-3 times at intervals of 15-20 days.
- Apply Isoprocarb (MIPC) group pesticide if banana beetle infestation occurs.
- In case of infestation of milli bugs on papaya trees, the infected leaves / stems including insects should be collected and destroyed at the beginning of the attack. If the infestation is severe, mix 5 gms of soapy water or Admire 200 ml at the rate of 0.25 ml per liter of water and spray 2-3 times in 15 days.
- Drainage out excess water from the land.
- Apply pesticides after current spell of rainfall.

Livestock

- Cattle should be fed with hay, raw grass and hand-made granular feed. Preserve paddy straw properly now.
- Vaccinate the cattle to prevent disease.

- Keep the cattle house dry and clean. Use mosquito nets or earthenware pots coils at room.
- Keep cattle safe place from lightning and rain.

Poultry

- Vaccinate the chickens regularly to prevent disease.
- Keep the poultry house dry and clean. Use mosquito nets or earthenware pots coils at room.

Fishery

- Now is the right time to stock fish. Prepare the pond.
- Catch all the old fish by drying the pond or applying Rotenon (25 g / decimal / ft). After 2/3 days apply 1 kg lime per decimal
- Apply fertilizer (200 g urea per decimal, 100 g TSP) and 3/4 days after lime application.
- 3/4 days after application of fertilizer, if the water colour is green, store the layer-based fish fry.

Realized Weather at different locations in the country

Rainfall for last 24 hours (till 06 AM on 25 August 2021) and maximum temperature on 24 August and minimum temperature on 25 August 2021 is given below:

Name of Divisions	Name of Stations	Rain fall (mm)	Max. Temp (°C)	Min. Temp (°C)	Name of Divisions	Name of Stations	Rain fall (mm)	Max. Temp (°C)	Min. Temp (°C)
Dhaka	Dhaka	09	34.2	25.7	Rajshahi	Rajshahi	00	35.0	27.0
	Tangail	09	33.4	25.8		Ishurdi	03	35.0	26.7
	Faridpur	17	33.9	25.7		Bogura	05	34.0	26.5
	Madaripur	01	34.0	27.0		Badalgachhi	20	33.8	26.4
	Gopalganj	02	33.3	26.8		Tarash	33	xx	25.5
	Nikli	43	33.0	25.4					
Mymensingh	Mymensingh	04	31.0	26.4	Rangpur	Rangpur	09	33.0	26.5
	Netrokona	<u>59</u>	29.0	25.5		Dinajpur	00	34.0	27.0
						Sayedpur	01	33.3	27.0
Chattogram	Chattogram	26	32.5	24.5		Tetulia	08	30.8	26.0
	Sandwip	04	32.1	27.2		Dimla	38	32.5	27.0
	Sitakunda	00	31.7	26.3		Rajarhat	35	32.0	25.5
	Rangamati	53	30.8	25.0	Khulna	Khulna	16	34.8	26.3
	Cumilla	01	32.6	26.2		Mongla	00	35.5	27.5
	Chandpur	00	<u>35.7</u>	28.0		Satkhira	00	32.8	27.5
	M.Court	00	33.4	27.6		Jashore	10	35.4	25.8
	Feni	09	32.2	25.5		Chuadanga	47	35.4	26.2
	Hatiya	00	33.5	28.0	Kumarkhali	11	33.0	25.5	
	Cox's Bazar	14	30.0	24.8	Barishal	Barishal	00	34.7	27.9
Kutubdia	43	32.5	26.1	Patuakhali		03	33.6	28.0	
Teknaf	30	30.9	xx	Khepupara		00	33.7	28.5	
Sylhet	Sylhet	13	29.2	25.0		Bhola	00	34.8	28.5
	Srimangal	28	33.7	<u>24.4</u>					

Highlights:


- Country average bright sunshine hour was 2.87 hours per day during last week.
- Country average free water loss during the previous week was 2.59 mm per day.





Weather Forecast Valid for 24 Hours commencing 06 PM Today:





Forecast: Light to moderate rain/thunder showers accompanied by temporary gusty wind is likely to occur at most places over Rangpur, Rajshahi, Mymensingh, Sylhet & Dhaka divisions and at many places over Chattogram, Khulna & Barishal divisions with moderately heavy to heavy falls at places over the country.

Temperature: Day and night temperature may fall slightly over the country.

Spatial Distribution of weather parameters for the Week ending 24 August 2021









Weather Forecast and Advisory for the Period of 23.08.2021 to 31.08.2021


Bright sunshine hour is likely to be from 3.00 to 4.00 hours per day during this week. Average free water loss is likely to be from 2.75 to 3.75 mm per day during this week.

- Light (04-10 mm/d) to moderate (11-22 mm/d) rain/thunder showers accompanied by temporary gusty wind is likely to occur at many places over Rajshahi, Rangpur, Mymensingh, Dhaka and Sylhet divisions and at a few places over Khulna, Barishal and Chattogram divisions with moderately heavy (23-43 mm/d) falls at places over the country during the 1st half of the period.
- Light (04-10 mm/d) to moderate (11-22 mm/d) rain/thunder showers accompanied by temporary gusty wind is likely to occur at many places over Khulna, Barishal and Chattogram divisions and at a few places over Dhaka, Rangpur, Rajshahi, Mymensingh and Sylhet divisions with moderately heavy (23-43 mm/d) falls at places over the country during the 2nd half of the period.
- Day and night temperature may rise slightly during the 1st half and it may falls slightly over the country during the 2nd half of the period.

District Wise Quantitative Weather Forecast for next Five days from 25 August-29 August 2021









Different Satellite Products over Bangladesh


NOAA/VIIRS BLENDED NDVI composite for the week No. 32 (06 August- 12 August) over Agricultural regions of Bangladesh



NOAA/ AVHRR BLENDED VCI composite for the week No. 32 (06 August- 12 August) over Agricultural regions of Bangladesh



NOAA/ AVHRR BLENDED TCI composite for the week No. 32 (06 August- 12 August) over Agricultural regions of Bangladesh



NOAA/ AVHRR BLENDED VHI composite for the week No. 32 (06 August- 12 August) over Agricultural regions of Bangladesh

