



# NATIONAL AGROMETEOROLOGICAL ADVISORY SERVICE BULLETIN

22 September 2021 (WEDNESDAY)  
[FOR THE PERIOD 22.09.2021 – 26.09.2021]



## DISCLAIMER

Under the Agro-meteorological Information System Development Project, Agromet Advisory Service Bulletins at National & Sixty Four (64) districts have been prepared on experimental basis. These bulletins are being displayed through the website of Department of Agricultural Extension for valuable comments/ suggestions from all concerned.

Please Contact: Farhana Hoque, Sabuj Roy | E-mail: [adamsdp@dae.gov.bd](mailto:adamsdp@dae.gov.bd)  
Phone No.: 02-55'028414, 02-55028418

## ISSUED BY

Agro-Meteorological Information Systems  
Development Project  
Department of Agricultural Extension (DAE)  
Ministry of Agriculture  
In collaboration with Bangladesh  
Meteorological Department

## **Special Agromet Advisories for CORONA virus (COVID-19) transmission in Bangladesh**

### **Agromet Advisories**

Farmers and all other farm managements are advised to follow the guidelines of Government of Bangladesh to avoid infection and social transmission of CORONA virus (COVID-19). Precautions and safety measures should be taken up to prevent the Carona virus spread. Simple measures include social distancing, maintaining personal hygiene by washing hands with soap, wearing of face mask, drink hot water, stay at home and cleaning of implements and machinery. Farmers should not work in a group; consult with a doctor in case of any symptom. Please stay at home does not visit the field until it is very necessary.

### **Salient Weather Conditions & Forecast**

As per the weather forecast of Bangladesh Meteorological Department (BMD) during next 24 hours' light to moderate rain/thunder showers accompanied by temporary gusty wind is likely to occur at many places over Khulna division; at a few places over Rajshahi & Barishal divisions and at one or two places over Rangpur, Dhaka Mymensingh, Chattogram & Sylhet divisions with moderately heavy falls at isolated places over southern part of the country. Day and night temperature may remain nearly unchanged over the country. As per the outlook issued by BMD for next 72 hours' rain/thunder showers may increase at the end period. During last four days, light rainfall to very heavy rainfall occurred many of the district of the country and as per the quantitative medium range weather forecast; many of the districts of the country are likely to receive light rainfall to heavy rainfall in the next five days. Detail Agromet advisory are given below:

#### **Aus:**

- Maintain the required water level in the land.
- Insecticides of carbosulfan group can be applied to control thrips, green leaf hopper and Rice Bug insects.
- Apply pesticides after current spell of rainfall.
- Drain out the water from the rice field 15 days before harvest.
- Collect mature crops after rains and keep them in a safe place quickly.

#### **Aman**

- Maintain the required water level in the land.
- In case of infestation of stem borer insects, apply 180-190 gm of Cartap group or 10 gm of Thiamethoxam + Chlorantraniliprol group of pesticides per bigha for control.
- In order to control Sheath Blight Disease, potash fertilizer should be divided into two equal installments, one part should be applied in the final cultivation of land preparation and the other part should be applied in combination with the last installment of urea fertilizer.
- In case of bacterial blight, in the early stages of the disease, mix 60 gm MOP, 60 gm thiovit in 10 liters of water and apply on 5 decimal of the land. If the disease occurs before the heading stage, 5 kg of potash fertilizer should be applied per bigha.

- In case of infestation of leaf-wrapping insects, apply 180-190 gm of Cartap group or 10 gm of Thiamethoxam + Chlorantraniliprole group of pesticides per bigha for control.
- Use light traps to control insects.
- Apply pesticides after current spell of rainfall.
- Drain out excess water from the land, if any.

### **Sugarcane**

- Sugarcane stems can be attacked by red rot disease. Infected trees should be removed from the land along with bushes as soon as the disease is seen. Water should be removed from the sugarcane field very quickly.
- To control the Sugarcane Stem Borer, remove the soil from the base of the sugarcane with a spade. Drain out accumulated water from the land. Old dried leaves should be removed from the trees, collected and burnt or buried under the ground. Egg parasitic hornet *Trichogramma Chilonis* should be released in the affected land at the rate of one gram per hectare per week (approximately 50,000). Apply Virtako 40 WG (Thiamethoxam + Chlorantraniliprole) to the infested sugarcane bush in the approved dose for control of stem borer insects with pesticides or Cartop granular pesticides such as Nokotap 6G cut the shallow drains on both sides of the sugarcane row and sprinkle in the drains then cover with soil or sprinkle at the base of the tree.
- Apply pesticides after current spell of rainfall.
- Drain out excess water from the land, if any.

### **Vegetables**

- Arrange for hand pollination of pumpkin vegetables. If beetles appear, kill them by hand in the morning and afternoon.
- In case of fly infestation in pumpkin, use pheromone and poison traps together. Pesticides of Alpha-Cypermethrin group can be used.
- In case of infestation of *eggplant* tip and fruit borer insects, cut and destroy the affected tip along with the insects. Insect reproduction is likely to be greatly reduced by using pheromone traps. Use spray of approval dose of insecticides, if necessary or recommended local organic insecticides may be used.
- In case of spider infestation, soak 1 kg half-broken neem seeds in 20 liters of water for 24 hours and spray the water (after chopping) on the underside of the leaves. If the infestation is severe, spray the spider mite Omit 57 EC at the rate of 2.0 ml per liter of water or Vertimek 1.8 EC at the rate of 12 ml per 10 liters of water.
- Apply pesticides after current spell of rainfall.
- Drain out excess water from the land, if any.

### **Horticultural crops**

- In case of spider infestation on the coconut tree, all the young caconut of the affected tree should be cut down and burnt. Infected trees and nearby young trees should be

sprayed with any spider repellent such as Omite 57 EC mixed in 1.5 ml of one liter of water.

- In case of symptoms of Sigatoka disease on banana leaves, 0.5 ml of Score or 2 g of Noin or Bavistin or 0.1 ml of Econazole / Folico should be mixed in every liter of water and sprayed 2-3 times at intervals of 15-20 days.
- Apply Isoprocarb (MIPC) group pesticide if banana beetle infestation occurs.
- In case of infestation of milli bugs on papaya trees, the infected leaves / stems including insects should be collected and destroyed at the beginning of the attack. If the infestation is severe, mix 5 gms of soapy water or Admire 200 ml at the rate of 0.25 ml per liter of water and spray 2-3 times in 15 days.
- Apply pesticides after current spell of rainfall.
- Drain out excess water from the land, if any.

#### **Livestock:**

- Cattle should be fed with hay, raw grass and hand-made granular feed.
- Vaccinate cattle regularly to prevent disease.
- Keep the cattle house dry and clean. Use mosquito nets or earthenware pots coils at room.
- Keep cattle safe place from lightning and rain.

#### **Poultry:**

- Vaccinate chickens regularly to prevent disease.
- Keep the poultry house dry and clean.
- Sprinkle potassium permanganate solution to remove odors around.

#### **Fishery**

- Large catchable fish in ponds sell in the fish Market.
- Re-reserve more than 20% of large size (6-8 kg) fish of the same species.

## Realized Weather at different locations in the country

Rainfall for last 24 hours (till 06 AM on 22 September 2021) and maximum temperature on 21 September and minimum temperature on 22 September 2021 is given below:

Name of Divisions	Name of Stations	Rain fall (mm)	Max. Temp (°C)	Min. Temp (°C)	Name of Divisions	Name of Stations	Rain fall (mm)	Max. Temp (°C)	Min. Temp (°C)
Dhaka	Dhaka	Trace	32.7	26.8	Rajshahi	Rajshahi	16	33.6	25.8
	Tangail	10	31.5	24.5		Ishurdi	03	31.3	25.8
	Faridpur	00	31.5	26.0		Bogura	12	31.5	25.5
	Madaripur	00	32.0	25.8		Badalgachhi	05	31.5	25.0
	Gopalganj	06	30.2	25.8		Tarash	06	31.0	26.0
	Nikli	05	32.5	25.5					
Mymensingh	Mymensingh	03	30.6	25.4	Rangpur	Rangpur	13	33.0	24.4
	Netrokona	01	30.5	25.4		Dinajpur	12	33.6	24.5
						Sayedpur	01	33.4	24.2
Chattogram	Chattogram	00	33.0	25.5		Tetulia	13	<b>35.5</b>	<b>23.1</b>
	Sandwip	00	31.7	25.5		Dimla	00	33.5	24.4
	Sitakunda	00	32.5	xx		Rajarhat	06	32.2	24.5
	Rangamati	00	32.5	24.8	Khulna	Khulna	09	30.5	25.5
	Cumilla	00	32.2	25.2		Mongla	19	29.0	25.0
	Chandpur	00	32.5	25.5		Satkhira	01	29.6	26.5
	M.Court	00	31.6	26.0		Jashore	Trace	31.4	25.6
	Feni	00	33.0	26.0		Chuadanga	Trace	31.5	25.7
	Hatiya	09	31.4	25.0		Kumarkhali	02	31.5	25.3
	Cox's Bazar	00	32.2	25.2	Barishal	Barishal	00	31.2	25.5
Kutubdia	00	31.8	25.5	Patuakhali		01	30.5	25.7	
Teknaf	00	32.0	28.0	Khepupara		09	29.4	25.5	
Sylhet	Sylhet	04	31.5	24.9	Bhola	<b>21</b>	31.0	25.5	
	Srimangal	00	31.5	24.7					

### Highlights:

- Country average bright sunshine hour was 3.82 hours per day during last week.
- Country average free water loss during the previous week was 2.93 mm per day.


### Weather Forecast Valid for 24 Hours commencing 06 PM Today:





**Forecast:** Light to moderate rain/thunder showers accompanied by temporary gusty wind is likely to occur at many places over Khulna division; at a few places over Rajshahi & Barishal divisions and at one or two places over Rangpur, Dhaka Mymensingh, Chattogram & Sylhet divisions with moderately heavy falls at isolated places over southern part of the country.

**Temperature:** Day and night temperature may remain nearly unchanged over the country.







Spatial Distribution of weather parameters for the Week ending 21 September 2021













### Weather Forecast and Advisory for the Period of 22.09.2021 to 30.09.2021


Bright sunshine hour is likely to be from 3.50 to 4.50 hours per day during this week. Average free water loss is likely to be from 3.00 to 4.00 mm per day during this week.

- Light (04-10 mm/d) to moderate (11-22 mm/d) rain/thunder showers accompanied by temporary gusty wind is likely to occur at a few places over Rajshahi, Rangpur, Dhaka, Khulna, Barishal, Chattogram, Mymensingh & Sylhet divisions with moderately heavy (23- 43 mm/d) falls at isolated places over the country during the 1st half of the period.
- Light (04-10 mm/d) to moderate (11-22 mm/d) rain/thunder showers accompanied by temporary gusty wind is likely to occur at many places over Dhaka, Khulna, Barishal & Chattogram divisions and at a few places over Rajshahi, Rangpur, Mymensingh & Sylhet divisions with moderately heavy (23-43 mm/d) to heavy (44-88 mm/d) falls at isolated places over the country during the 2nd half of the period.
- Day and night temperature may remain nearly unchanged during the 1st half and it may fall slightly over the country during the 2nd half of the period.

### **District Wise Quantitative Weather Forecast for next Five days from 22 Sept.-26 September 2021**











## Different Satellite Products over Bangladesh


NOAA/VIIRS BLENDED NDVI composite for the week No. 36 (03 September-09 September) over Agricultural regions of Bangladesh



NOAA/ AVHRR BLENDED VCI composite for the week No. 36 (03 September- 09 September) over Agricultural regions of Bangladesh



NOAA/ AVHRR BLENDED TCI composite for the week No. 36 (03 September-09 September) over Agricultural regions of Bangladesh



NOAA/ AVHRR BLENDED VHI composite for the week No. 36 (03 September- 09 September) over Agricultural regions of Bangladesh

