Faridpur Region		Rabi season Garlic: Thrips insect												Life Span 140-150 day										
Weather warning	Months	November					December				January				February				March					
	Max. Temp. (oC)						30.84 °C, hot and dry weather increases Thrips population also associated with lack of rainfall																	
	Min. Temp. (oC)	17.5 °C, Develop slowly during winter or lower temperatures																						
	RH Max (%)	78.62%, High relative humidity reduces Thrips																						
	RH Min (%)	47.6%, total life duration shortest in high temperature and low humidity																						
	Rainfall (mm)	Heavy rainfall reduces population																						
	Std.Week/Normal	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8	9	10	11	12	13	
Weekly normal weather	Rainfall (mm)	11.5	11.0	8.0	3.0	4.5	0.5	3.5	0.5	2.0	0.5	0.5	0.5	1.5	3.5	5.5	5.0	7.5	5.0	6.5	7.0	12.5	18.5	
	Max. Temp. (oC)	30.9	30.4	29.7	29.2	28.0	27.5	26.5	25.5	25.1	24.3	24.4	24.8	25.4	26.5	27.4	28.5	29.5	30.7	31.8	32.9	33.8	33.4	
	Min. Temp. (oC)	22.0	20.7	19.1	18.2	16.4	15.2	14.7	13.7	12.9	12.2	12.0	12.1	12.1	13.1	13.9	15.5	16.0	17.3	18.5	19.5	21.2	21.8	
	RHmax (%)	95.2	95.4	95.5	95.8	95.6	95.6	96.1	96.2	96.0	95.7	95.4	95.5	95.2	94.7	94.7	64.1	93.8	93.4	92.7	92.4	92.9	93.2	
	RHMin (%)	57.6	55.5	53.2	51.2	49.9	50.5	51.4	51.3	51.0	52.2	50.9	48.9	46.3	46.2	43.3	43.3	41.5	39.5	38.8	38.4	42.1	46.7	
	SShr (hrs)	48.5	51.0	49.5	50.5	51.0	47.0	44.5	40.5	43.0	41.5	39.5	44.5	50.0	49.0	52.0	52.0	53.0	54.5	53.0	54.5	51.5	50.5	
Stage		Germination and Seedling Planting					Vegetative Growth					Bulb formation						Harvesting						
Inse	ect Life Cycle						Eggs (2 to 6 days)	Larvae (5 to 7 days)		Pupa 10 days)		to 30 t	gs (2 o 6 ays)	Larvae (5 to 7 days)	Pu (3 1 da	to 0	Adults (28 to 30 days)							

Crop insect weather calendar of the Rabi season Gurlic (Thrips insect): Faridpur Region (Districts: Faridpur, Madaripur, Sariatpur, Gopalgonj, Rajbari) Bangladesh