



Special Post-Cyclone 'YAAS' Agromet Advisories

(Applicable for Satkhira, Khulna, Bagerhat, Pirojpur, Barguna, Jhalokati, Barishal, Patuakhali, Bhola, Naokhali, Feni, Lakshmipur, Chandpur, Chattogram and Cox'sBazar)

Published date: 28 May 2021

Agromet advisories are given below:

- Drain out excess water immediately from boro paddy field as well as other standing crops.
- Harvest the matured boro paddy and other crops after draining out the excess water.
- Immediately sundry the wet boro paddy and store in safe and dry place.
- If Aus paddy has not been sown, sow immediately after draining out water from the field.
- After 5-7 days of water drainage from Aus paddy field, filling up the gaps and then apply potash fertilizer as top dress. After 3-5 days of it apply also urea fertilizer. Urea fertilizer should not be applied if the paddy is in early tillering stage. After 5-7 days, if necessary, urea fertilizer should be applied by breaking the tillers and filling the gap. At this time, it is better to use additional 5 kg of potash fertilizer per bigha in all types of Aus paddy fields.
- Drain out water, if any, immediately from jute field. Application of fertiliser to jute field should be avoided now.
- After drying the jute land, undertake weeding operation with a hoe. In case of disease incidences, apply recommended fungicides.
- Immediately raise the lodged banana and other horticultural crops by mechanical support.
- Start sowing of vegetables and other crops under optimum soil moisture condition.
- Thinning, weeding and other intercultural operations should be done after draining out excess water from the crop field.
- Apply the required fertilizers and pesticides after draining out excess water from the crop field.

- Before water drained out from the field, vegetables like Sponge gourd, Dhundal, Bitter gourd, Pumpkin etc. can be planted in the backyard and road side.
- Keep cattle and poultry on a high land & clean area. Also provide foods rich in vitamins and minerals.
- Repair all the submerged ponds as soon as the water drain out.
- On a sunny day, lime should be applied to the ponds at the rate of 250-650 g / per decimal depending on the quantity of fish. Three days after lime application, urea at the rate of 80-100 g / per decimal and TSP at the rate of 40-50 g / per decimal should be applied on sunny days.
- Check whether the fish has come out of the pond by pulling the net. In the ponds where the fish have gone out, a little bigger fry need to be restocked.
- Carry out all the farm operations by maintaining social distancing.



(Dr. Md. Shah Kamal Khan)

Project Director

Cell no. +8801712184274

Telephone: +88-0255028422